

U.S. PATENT APPLICATION

Inventor(s): Jane YE
Chunhua YAN
Valentina DI FRANCESCO
Ellen M. BEASLEY

Invention: ISOLATED HUMAN RAS-LIKE PROTEINS, NUCLEIC ACID
MOLECULES ENCODING THESE HUMAN RAS-LIKE PROTEINS, AND
USES THEREOF

CELERA GENOMICS CORPORATION.
45 WEST GUDE DR., C2-4#20
ROCKVILLE, MD 20850
(240) 453-3067
Fax (240)-453-3084

SPECIFICATION

ISOLATED HUMAN RAS-LIKE PROTEINS, NUCLEIC ACID MOLECULES ENCODING THESE HUMAN RAS-LIKE PROTEINS, AND USES THEREOF

FIELD OF THE INVENTION

The present invention is in the field of Ras-like proteins that are related to the Rab subfamily, recombinant DNA molecules and protein production. The present invention specifically provides novel Ras-like protein polypeptides and proteins and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

BACKGROUND OF THE INVENTION

Ras-like proteins, particularly members of the Rab subfamilies, are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown members of this subfamily of Ras-like proteins. The present invention advances the state of the art by providing a previously unidentified human Ras-like proteins that have homology to members of the Rab subfamilies.

Ras protein

Ras proteins are small regulatory GTP-binding proteins, or small G proteins, which belong to the Ras protein superfamily. They are monomeric GTPases, but their GTPase activity is very slow (less than one GTP molecule per minute).

Ras proteins are key relays in the signal-transducing cascade induced by the binding of a ligand to specific receptors such as receptor tyrosine kinases (RTKs), since they trigger the MAP kinase cascade. The ligand can be a growth factor (epidermal growth factor (EGF), platelet-derived growth factor (PDGF), insulin, an interleukin (IL), granulocyte colony-stimulating factor (G-CSF), granulocyte/macrophage colony-stimulating factor (GM-CSF).

Ras proteins contain sequences highly conserved during evolution. Their tertiary structure includes ten loops connecting six strands of beta-sheet and five alpha helices.

In mammals, there are four Ras proteins, which are encoded by Ha-ras, N-ras, Ki-rasA and Ki-rasB genes. They are composed of about 170 residues and have a relative molecular mass of 21 kD. Ras proteins contain covalently attached modified lipids allowing these proteins to bind to the plasma membrane. Ha-Ras has a C-terminal farnesyl group, a C-terminal palmitoyl group and a N-terminal myristoyl group. In Ki-Ras(B), a C-terminal polylysine domain replaces the palmitoyl group.

Ras proteins alternate between an inactive form bound to GDP and an active form bound to GTP. Their activation results from reactions induced by a guanine nucleotide-exchange factor (GEF). Their inactivation results from reactions catalyzed by a GTPase-activating protein (GAP).

When a Ras protein is activated by a GEF such as a Sos protein, the N-terminal region of a serine/threonine kinase, called "Raf protein", can bind to Ras protein. The C-terminal region of the activated Raf thus formed binds to another protein, MEK, and phosphorylates it on both specific tyrosine and serine residues. Active MEK phosphorylates and activates, in turn, a MAP kinase (ERK1 or ERK2), which is also a serine/threonine kinase. This phosphorylation occurs on both specific tyrosine and threonine residues of MAP kinase.

MAP kinase phosphorylates many different proteins, especially nuclear transcription factors (TFs) that regulate expression of many genes during cell proliferation and differentiation.

Recent researches suggest that, in mammals, phosphatidyl inositol 3'-kinase (PI3-kinase) might be a target of Ras protein, instead of Raf protein. In certain mutations, the translation of ras genes may produce oncogenic Ras proteins.

Ras-like protein

Guanine nucleotide-binding proteins (GTP-binding proteins, or G proteins) participate in a wide range of regulatory functions including metabolism, growth, differentiation, signal transduction, cytoskeletal organization, and intracellular vesicle transport and secretion. These proteins control diverse sets of regulatory pathways in

response to hormones, growth factors, neuromodulators, or other signaling molecules. When these molecules bind to transmembrane receptors, signals are propagated to effector molecules by intracellular signal transducing proteins. Many of these signal-transducing proteins are members of the Ras superfamily.

The Ras superfamily is a class of low molecular weight (LMW) GTP-binding proteins that consist of 21-30 kDa polypeptides. These proteins regulate cell growth, cell cycle control, protein secretion, and intracellular vesicle interaction. In particular, the LMW GTP-binding proteins activate cellular proteins by transducing mitogenic signals involved in various cell functions in response to extracellular signals from receptors (Tavitian, A. (1995) C. R. Seances Soc. Biol. Fil. 189:7-12). During this process, the hydrolysis of GTP acts as an energy source as well as an on-off switch for the GTPase activity of the LMW GTP-binding proteins.

The Ras superfamily is comprised of five subfamilies: Ras, Rho, Ran, Rab, and ADP-ribosylation factor (ARF). Specifically, Ras genes are essential in the control of cell proliferation. Mutations in Ras genes have been associated with cancer. Rho proteins control signal transduction in the process of linking receptors of growth factors to actin polymerization that is necessary for cell division. Rab proteins control the translocation of vesicles to and from membranes for protein localization, protein processing, and secretion. Ran proteins are localized to the cell nucleus and play a key role in nuclear protein import, control of DNA synthesis, and cell-cycle progression. ARF and ARF-like proteins participate in a wide variety of cellular functions including vesicle trafficking, exocrine secretion, regulation of phospholipase activity, and endocytosis.

Despite their sequence variations, all five subfamilies of the Ras superfamily share conserved structural features. Four conserved sequence regions (motifs I-IV) have been studied in the LMW GTP-binding proteins. Motif I is the most variable but has the conserved sequence, GXXXXGK. The lysine residue is essential in interacting with the .beta.- and .gamma.-phosphates of GTP. Motif II, III, and IV contain highly conserved sequences of DTAGQ, NKXD, and EXSAX, respectively. Specifically, Motif II regulates the binding of gamma-phosphate of GTP; Motif III regulates the binding of GTP; and Motif IV regulates the guanine base of GTP. Most of the membrane-bound LMW GTP-binding proteins generally require a carboxy terminal isoprenyl group for membrane

association and biological activity. The isoprenyl group is added posttranslationally through recognition of a terminal cysteine residue alone or a terminal cysteine-aliphatic amino acid-aliphatic amino acid-any amino acid (CAAX) motif. Additional membrane-binding energy is often provided by either internal palmitoylation or a carboxy terminal cluster of basic amino acids. The LMW GTP-binding proteins also have a variable effector region, located between motifs I and II, which is characterized as the interaction site for guanine nucleotide exchange factors (GEFs) or GTPase-activating proteins (GAPs). GEFs induce the release of GDP from the active form of the G protein, whereas GAPs interact with the inactive form by stimulating the GTPase activity of the G protein.

The ARF subfamily has at least 15 distinct members encompassing both ARF and ARF-like proteins. ARF proteins identified to date exhibit high structural similarity and ADP-ribosylation enhancing activity. In contrast, several ARF-like proteins lack ADP-ribosylation enhancing activity and bind GTP differently. An example of ARF-like proteins is a rat protein, ARL184. ARL184 has been shown to have a molecular weight of 22 kDa and four functional GTP-binding sites (Icard-Liepkalns, C. et al. (1997) *Eur. J. Biochem.* 246: 388-393). ARL184 is active in both the cytosol and the Golgi apparatus and is closely associated with acetylcholine release, suggesting that ARL184 is a potential regulatory protein associated with Ca^{2+} -dependent release of acetylcholine.

A number of Rho GTP-binding proteins have been identified in plasma membrane and cytoplasm. These include RhoA, B and C, and D, rhoG, rac 1 and 2, G25K-A and B, and TC10 (Hall, A. et al. (1993) *Philos. Trans. R. Soc. Lond. (Biol.)* 340:267-271). All Rho proteins have a CAAX motif that binds a prenyl group and either a palmitoylation site or a basic amino acid-rich region, suggesting their role in membrane-associated functions. In particular, RhoD is a protein that functions in early endosome motility and distribution by inducing rearrangement of actin cytoskeleton and cell surface (Murphy, C. et al. (1996) *Nature* 384:427-432). During cell adhesion, the Rho proteins are essential for triggering focal complex assembly and integrin-dependent signal transduction (Hotchin, N. A. and Hall, A. (1995) *J. Cell Biol.* 131:1857-1865).

The Ras subfamily proteins already indicated supra are essential in transducing signals from receptor tyrosine kinases (RTKs) to a series of serine/threonine kinases

which control cell growth and differentiation. Mutant Ras proteins, which bind but cannot hydrolyze GTP, are permanently activated and cause continuous cell proliferation or cancer. TC21, a Ras-like protein, is found to be highly expressed in a human teratocarcinoma cell line (Drivas, G. T. et al. (1990) *Mol. Cell. Biol.* 10: 1793-1798). Rin and Rit are characterized as membrane-binding, Ras-like proteins without the lipid-binding CAAX motif and carboxy terminal cysteine (Lee, C.-H. J. et al. (1996) *J. Neurosci.* 16: 6784-6794). Further, Rin is shown to localize in neurons and have calcium-dependant calmodulin-binding activity.

Rab proteins

The novel human protein, and encoding gene, provided by the present invention is related to the Rab subfamily of Ras proteins and shows a high degree of similarity to Rab15. Furthermore, the protein of the present invention may be a novel alternative splice form of a protein provided in WO200058473; specifically, the protein of the present invention differs from the art-known protein of WO200058473 in exon 6 (see amino acid sequence alignment in Figure 2).

Rab15 may act, together with Rab3A, to regulate synaptic vesicle membrane flow within nerve terminals, thereby regulating neurotransmitter release. Rab15 and Rab3A are low molecular weight GTP-binding proteins. Rab proteins are generally comprised of four conserved structural domains necessary for GTP binding, as well as additional domains for membrane localization and effector protein interactions. Rab15 is expressed primarily in neural tissues such as the brain and is localized to synaptic vesicles (Elferink *et al.*, *J. Biol. Chem.* 267 (9), 5768-5775 (1992)).

Due to their importance in neural physiology, particularly in regulating neurotransmitter release, novel human Rab proteins/genes, such as provided by the present invention, are valuable as potential targets for the development of therapeutics to treat neurological disorders. Furthermore, SNPs in Rab genes, such as provided by the present invention, are valuable markers for the diagnosis, prognosis, prevention, and/or treatment of neurological disorders.

Using the information provided by the present invention, reagents such as probes/primers for detecting the SNPs or the expression of the protein/gene provided

herein may be readily developed and, if desired, incorporated into kit formats such as nucleic acid arrays, primer extension reactions coupled with mass spec detection (for SNP detection), or TaqMan PCR assays.

The discovery of new human Ras-like proteins and the polynucleotides that encode them satisfies a need in the art by providing new compositions that are useful in the diagnosis, prevention, and treatment of inflammation and disorders associated with cell proliferation and apoptosis.

SUMMARY OF THE INVENTION

The present invention is based in part on the identification of amino acid sequences of human Ras-like protein polypeptides and proteins that are related to the Rab Ras-like protein subfamily, as well as allelic variants and other mammalian orthologs thereof. These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and serve as targets for the development of human therapeutic agents that modulate Ras-like protein activity in cells and tissues that express the Ras-like protein. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors.

DESCRIPTION OF THE FIGURE SHEETS

FIGURE 1 provides the nucleotide sequence of a cDNA molecule that encodes the Ras-like protein of the present invention. (SEQ ID NO:1) In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of inventions based on this molecular sequence. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors.

FIGURE 2 provides the predicted amino acid sequence of the Ras-like protein of the present invention. (SEQ ID NO:2) In addition structure and functional information such as protein family, function, and modification sites is provided where available,

allowing one to readily determine specific uses of inventions based on this molecular sequence.

FIGURE 3 provides genomic sequences that span the gene encoding the Ras-like protein of the present invention. (SEQ ID NO:3) In addition structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence. As illustrated in Figure 3, SNPs were identified at 20 different nucleotide positions.

DETAILED DESCRIPTION OF THE INVENTION

General Description

The present invention is based on the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a Ras-like protein or part of a Ras-like protein and are related to the Rab subfamily. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human Ras-like protein polypeptides that are related to the Rab subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these Ras-like protein polypeptide, nucleic acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the Ras-like protein of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known Ras-like proteins of the Rab subfamily and the expression pattern observed. Experimental data as provided in

Figure 1 indicates expression in the brain and in tumors. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known Rab family or subfamily of Ras-like proteins.

Specific Embodiments

Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the Ras-like protein family and are related to the Rab subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figure 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the Ras-like proteins or peptides of the present invention, Ras-like proteins or peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprise the amino acid sequences of the Ras-like protein polypeptide disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is

that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components.

In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the Ras-like protein polypeptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated Ras-like protein polypeptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. For example, a nucleic acid molecule encoding the Ras-like protein polypeptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence is the final amino acid sequence of the protein.

The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the Ras-like protein polypeptide of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The Ras-like protein polypeptides of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a Ras-like protein polypeptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the Ras-like protein polypeptide. "Operatively linked" indicates that the Ras-like protein polypeptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the Ras-like protein polypeptide.

In some uses, the fusion protein does not affect the activity of the Ras-like protein polypeptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins,

particularly poly-His fusions, can facilitate the purification of recombinant Ras-like protein polypeptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A Ras-like protein polypeptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the Ras-like protein polypeptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the peptides of the present invention, such as naturally occurring mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art know techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the Ras-like protein polypeptides of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family, and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix

and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, word length = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, word length = 3, to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the Ras-like protein polypeptides of the present invention as well as being encoded by the same genetic locus as the Ras-like protein polypeptide provided herein. The gene encoding the novel Ras-like protein of the present invention is located on a genome component that has been mapped to human chromosome 14 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data.

Allelic variants of a Ras-like protein polypeptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the Ras-like protein polypeptide as well as being encoded by the same genetic locus as the Ras-like protein polypeptide provided herein. Genetic locus can readily be determined based on the genomic information provided in Figure 3, such as the genomic

sequence mapped to the reference human. The gene encoding the novel Ras-like protein of the present invention is located on a genome component that has been mapped to human chromosome 14 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data. As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a Ras-like protein polypeptide encoding nucleic acid molecule under stringent conditions as more fully described below.

Figure 3 provides information on SNPs that have been found in the gene encoding the Ras-like protein of the present invention. SNPs were identified at 20 different nucleotide positions. Some of these SNPs that are located outside the ORF and in introns may affect gene transcription.

Paralogs of a Ras-like protein polypeptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the Ras-like protein polypeptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 40-50%, 50-60%, and more typically at least about 60-70% or more homologous through a given region or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a Ras-like protein polypeptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a Ras-like protein polypeptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the Ras-like protein polypeptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a Ras-like protein polypeptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the Ras-like protein polypeptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the Ras-like protein polypeptide. For example, one class of substitutions is conserved amino acid substitutions. Such substitutions are those that substitute a given amino acid in a Ras-like protein polypeptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg, replacements among the aromatic residues Phe, Tyr, and the like. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant Ras-like protein polypeptides can be fully functional or can lack function in one or more activities. Fully functional variants typically contain only conservative variations or variations in non-critical residues or in non-critical regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallography, nuclear magnetic resonance, or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

The present invention further provides fragments of the Ras-like protein polypeptides, in addition to proteins and peptides that comprise and consist of such

fragments. Particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that have been disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16 or more contiguous amino acid residues from a Ras-like protein polypeptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the Ras-like protein polypeptide, or can be chosen for the ability to perform a function, e.g., act as an immunogen. Particularly important fragments are biologically active fragments, peptides that are, for example about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the Ras-like protein polypeptide, e.g., active site. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known and readily available to those of skill in the art (e.g., PROSITE, HMMer, eMOTIF, etc.). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in Ras-like protein polypeptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation,

racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the Ras-like protein polypeptides of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature Ras-like protein polypeptide is fused with another compound, such as a compound to increase the half-life of the Ras-like protein polypeptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature Ras-like protein polypeptide, such as a leader or secretory sequence or a sequence for purification of the mature Ras-like protein polypeptide, or a pro-protein sequence.

Protein/Peptide Uses

The proteins of the present invention can be used in assays to determine the biological activity of the protein, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its ligand or receptor) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the binding partner so as to develop a system to identify inhibitors of the binding interaction. Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, Ras-like proteins isolated from humans and their human/mammalian orthologs serve as targets for identifying agents for use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the Ras-like protein. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain. A large percentage of pharmaceutical agents are being developed that modulate the activity of Ras-like proteins, particularly members of the Rab subfamily (see Background of the Invention). The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the

expression information provided in Figure 1. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. Such uses can readily be determined using the information provided herein, that which is known in the art, and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to Ras-like proteins that are related to members of the Rab subfamily. Such assays involve any of the known Ras-like protein functions or activities or properties useful for diagnosis and treatment of Ras-like protein-related conditions that are specific for the subfamily of Ras-like proteins that the one of the present invention belongs to, particularly in cells and tissues that express the Ras-like protein. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain.

The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the Ras-like protein, as a biopsy or expanded in cell culture. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the Ras-like protein.

The polypeptides can be used to identify compounds that modulate Ras-like protein activity. Both the Ras-like protein of the present invention and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the Ras-like protein. These compounds can be further screened against a functional Ras-like protein to determine the effect of the compound on the Ras-like protein activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the Ras-like protein to a desired degree.

Therefore, in one embodiment, Rab or a fragment or derivative thereof may be administered to a subject to prevent or treat a disorder associated with an increase in apoptosis. Such disorders include, but are not limited to, AIDS and other infectious or

genetic immunodeficiencies, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, and cerebellar degeneration, myelodysplastic syndromes such as aplastic anemia, ischemic injuries such as myocardial infarction, stroke, and reperfusion injury, toxin-induced diseases such as alcohol-induced liver damage, cirrhosis, and lathyrism, wasting diseases such as cachexia, viral infections such as those caused by hepatitis B and C, and osteoporosis.

In another embodiment, a pharmaceutical composition comprising Rab may be administered to a subject to prevent or treat a disorder associated with increased apoptosis including, but not limited to, those listed above.

In still another embodiment, an agonist which is specific for Rab may be administered to prevent or treat a disorder associated with increased apoptosis including, but not limited to, those listed above.

In a further embodiment, a vector capable of expressing Rab, or a fragment or a derivative thereof, may be used to prevent or treat a disorder associated with increased apoptosis including, but not limited to, those listed above.

In cancer, where Rab promotes cell proliferation, it is desirable to decrease its activity. Therefore, in one embodiment, an antagonist of Rab may be administered to a subject to prevent or treat cancer including, but not limited to, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, and teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. In one aspect, an antibody specific for Rab may be used directly as an antagonist, or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express Rab.

In another embodiment, a vector expressing the complement of the polynucleotide encoding Rab may be administered to a subject to prevent or treat a cancer including, but not limited to, the types of cancer listed above.

In inflammation, where Rab promotes cell proliferation, it is desirable to decrease its activity. Therefore, in one embodiment, an antagonist of Rab may be administered to a subject to prevent or treat an inflammation. Disorders associated with inflammation include,

but are not limited to, Addison's disease, adult respiratory distress syndrome, allergies, anemia, asthma, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, ulcerative colitis, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, atrophic gastritis, glomerulonephritis, gout, Graves' disease, hypereosinophilia, irritable bowel syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, rheumatoid arthritis, scleroderma, Sjogren's syndrome, and autoimmune thyroiditis; complications of cancer, hemodialysis, extracorporeal circulation; viral, bacterial, fungal, parasitic, protozoal, and helminthic infections and trauma. In one aspect, an antibody specific for Rab may be used directly as an antagonist, or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express Rab.

Further, the Ras-like protein polypeptides can be used to screen a compound for the ability to stimulate or inhibit interaction between the Ras-like protein and a molecule that normally interacts with the Ras-like protein, e.g. a ligand or a component of the signal pathway that the Ras-like protein normally interacts. Such assays typically include the steps of combining the Ras-like protein with a candidate compound under conditions that allow the Ras-like protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the Ras-like protein and the target, such as any of the associated effects of signal transduction.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries). (Hodgson, *Bio/technology*, 1992, Sept 10(9);973-80).

One candidate compound is a soluble fragment of the Ras-like protein that competes for ligand binding. Other candidate compounds include mutant Ras-like proteins or appropriate fragments containing mutations that affect Ras-like protein function and thus compete for ligand. Accordingly, a fragment that competes for ligand, for example with a higher affinity, or a fragment that binds ligand but does not allow release, is within the scope of the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) Ras-like protein activity. The assays typically involve an assay of events in the Ras-like protein mediated signal transduction pathway that indicate Ras-like protein activity. Thus, the phosphorylation of a protein/ligand target, the expression of genes that are up- or down-regulated in response to the Ras-like protein dependent signal cascade can be assayed. In one embodiment, the regulatory region of such genes can be operably linked to a marker that is easily detectable, such as luciferase. Alternatively, phosphorylation of the Ras-like protein, or a Ras-like protein target, could also be measured.

Any of the biological or biochemical functions mediated by the Ras-like protein can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art.

Binding and/or activating compounds can also be screened by using chimeric Ras-like proteins in which any of the protein's domains, or parts thereof, can be replaced by heterologous domains or subregions. Accordingly, a different set of signal transduction components is available as an end-point assay for activation. This allows for assays to be performed in other than the specific host cell from which the Ras-like protein is derived.

The Ras-like protein polypeptide of the present invention is also useful in competition binding assays in methods designed to discover compounds that interact with the Ras-like protein. Thus, a compound is exposed to a Ras-like protein polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble Ras-like protein polypeptide is also added to the mixture. If the test compound interacts with the soluble Ras-like protein polypeptide, it decreases the amount of complex

formed or activity from the Ras-like protein target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the Ras-like protein. Thus, the soluble polypeptide that competes with the target Ras-like protein region is designed to contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the Ras-like protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase/15625 fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., ^{35}S -labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of Ras-like protein-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin with techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a Ras-like protein-binding protein and a candidate compound are incubated in the Ras-like protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the Ras-like protein target molecule, or which are reactive with Ras-like protein and compete with the target molecule, as well as enzyme-

linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the Ras-like proteins of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal/insect model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of Ras-like protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the Ras-like protein associated pathway, by treating cells that express the Ras-like protein. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. These methods of treatment include the steps of administering the modulators of protein activity in a pharmaceutical composition as described herein, to a subject in need of such treatment.

In yet another aspect of the invention, the Ras-like proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos et al., *Cell* 72:223-232 (1993); Madura et al., *J. Biol. Chem.* 268:12046-12054 (1993); Bartel et al., *Biotechniques* 14:920-924 (1993); Iwabuchi et al., *Oncogene* 8:1693-1696 (1993); and Brent WO94/10300), to identify other proteins that bind to or interact with the Ras-like protein and are involved in Ras-like protein activity. Such Ras-like protein-binding proteins are also likely to be involved in the propagation of signals by the Ras-like proteins or Ras-like protein targets as, for example, downstream elements of a Ras-like protein-mediated signaling pathway, e.g., a pain signaling pathway. Alternatively, such Ras-like protein-binding proteins are likely to be Ras-like protein inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a Ras-like protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait"

and the "prey" proteins are able to interact, *in vivo*, forming a Ras-like protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the Ras-like protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a Ras-like protein modulating agent, an antisense Ras-like protein nucleic acid molecule, a Ras-like protein-specific antibody, or a Ras-like protein-binding partner) can be used in an animal or insect model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or insect model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The Ras-like proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to a disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. The method involves contacting a biological sample with a compound capable of interacting with the receptor protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells, and fluids present within a subject.

The peptides also are useful to provide a target for diagnosing a disease or predisposition to a disease mediated by the peptide. Accordingly, the invention provides

methods for detecting the presence, or levels of, the protein in a cell, tissue, or organism. The method involves contacting a biological sample with a compound capable of interacting with the receptor protein such that the interaction can be detected.

The peptides of the present invention also provide targets for diagnosing active disease, or predisposition to a disease, in a patient having a variant peptide. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in translation of an aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered receptor activity in cell-based or cell-free assay, alteration in ligand or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

In vitro techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence using a detection reagents, such as an antibody or protein binding agent.. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11) :983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both

the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the receptor protein in which one or more of the receptor functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other ligand-binding regions that are more or less active in ligand binding, and receptor activation. Accordingly, ligand dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. Accordingly, methods for treatment include the use of the Ras-like protein or fragments.

Antibodies

The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments thereof. As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')₂, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, Antibodies, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the Ras-like proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or receptor/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2)..

Detection of an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or

acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S , or ^3H .

Antibody Uses

The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues in an organism and over the course of normal development. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or abnormal expression during development. Antibody detection of circulating fragments of the full-length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data as provided in

Figure 1 indicates expression in the brain and in tumors. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the Ras-like protein to a binding partner such as a substrate. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in the sample with a standard; and instructions for use.

Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a Ras-like protein polypeptide of the present invention. Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the

Ras-like protein polypeptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule. The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript

sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprises several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, humans genomic sequence (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

Full-length genes may be cloned from known sequence using any one of a number of methods known in the art. For example, a method which employs XL-PCR (Perkin-Elmer, Foster City, Calif.) to amplify long pieces of DNA may be used. Other methods for obtaining full-length sequences are well known in the art.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when

the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life, or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the Ras-like protein polypeptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding, and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form of DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides nucleic acid molecules that encode fragments of the peptides of the present invention and that encode obvious variants of the Ras-like proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or whole organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions, and/or insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules provided in the Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences, and gene termination sequences. Such fragments are useful in controlling heterologous gene expression and in developing screens to identify gene-modulating agents.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could be at least 30, 40, 50, 100, 250, or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope-bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50, or more consecutive nucleotides.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. The gene encoding the novel Ras-like protein of the present invention is located on a genome component that has been mapped to human chromosome 14 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data.

Figure 3 provides information on SNPs that have been found in the gene encoding the Ras-like protein of the present invention. SNPs were identified at 20 different nucleotide positions. Some of these SNPs that are located outside the ORF and in introns may affect gene transcription.

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions can be such that sequences at least about 60%, at least about 70%, or at least about 80% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Examples of moderate to low stringency hybridization conditions are well known in the art.

Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2. As illustrated in Figure 3, SNPs were identified at 20 different nucleotide positions.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as those, which may encompass fragments disclosed prior to the present invention.

The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid

molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

The nucleic acid molecules are also useful as probes for determining the chromosomal positions of the nucleic acid molecules by means of *in situ* hybridization methods. The gene encoding the novel Ras-like protein of the present invention is located on a genome component that has been mapped to human chromosome 14 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data.

The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides. Moreover, the nucleic acid molecules are useful for constructing transgenic animals wherein a homolog of the nucleic acid molecule has been “knocked-out” of the animal’s genome.

The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form, and distribution of nucleic acid expression. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain. Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA

or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in Ras-like protein expression relative to normal results.

In vitro techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA include Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a Ras-like protein, such as by measuring a level of a receptor-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a receptor gene has been mutated. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate Ras-like protein nucleic acid expression.

The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the Ras-like protein gene, particularly biological and pathological processes that are mediated by the Ras-like protein in cells and tissues that express it. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors. The method typically includes assaying the ability of the compound to modulate the expression of the Ras-like protein nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired Ras-like protein nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the Ras-like protein nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

The assay for Ras-like protein nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the Ras-like protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of Ras-like protein gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of Ras-like protein mRNA in the presence of the candidate compound is compared to the level of expression of Ras-like protein mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate Ras-like protein nucleic acid expression in cells and tissues that express the Ras-like protein. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) of nucleic acid expression.

Alternatively, a modulator for Ras-like protein nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the Ras-like protein nucleic acid expression in the cells and tissues that express the protein. Experimental data as provided in Figure 1 indicates expression in the brain and in tumors.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the Ras-like protein gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the

compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in Ras-like protein nucleic acid, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in Ras-like protein genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally occurring genetic mutations in the Ras-like protein gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns, or changes in gene copy number, such as amplification. Detection of a mutated form of the Ras-like protein gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a Ras-like protein.

Individuals carrying mutations in the Ras-like protein gene can be detected at the nucleic acid level by a variety of techniques. Figure 3 provides information on SNPs that have been found in the gene encoding the Ras-like protein of the present invention. SNPs were identified at 20 different nucleotide positions. Some of these SNPs that are located outside the ORF and in introns may affect gene transcription. The gene encoding the novel Ras-like protein of the present invention is located on a genome component that has been mapped to human chromosome 14 (as indicated in Figure 3), which is supported by multiple lines of evidence, such as STS and BAC map data. Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the

gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a Ras-like protein gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant Ras-like protein gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., *Biotechniques* 19:448 (1995)), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79

(1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include, selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship). Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the Ras-like protein gene in an individual in order to select an appropriate compound or dosage regimen for treatment. Figure 3 provides information on SNPs that have been found in the gene encoding the Ras-like protein of the present invention. SNPs were identified at 20 different nucleotide positions. Some of these SNPs that are located outside the ORF and in introns may affect gene transcription.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control Ras-like protein gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of Ras-like protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into Ras-like protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of Ras-like protein nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired Ras-like protein nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of

the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the Ras-like protein, such as ligand binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in Ras-like protein gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired Ras-like protein to treat the individual.

The invention also encompasses kits for detecting the presence of a Ras-like protein nucleic acid in a biological sample. Experimental data as provided in Figure 1 indicates that Ras-like proteins of the present invention are expressed in the brain and in tumors, as indicated by virtual northern blot analysis. In addition, PCR-based tissue screening panels indicate expression in fetal brain. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting Ras-like protein nucleic acid in a biological sample; means for determining the amount of Ras-like protein nucleic acid in the sample; and means for comparing the amount of Ras-like protein nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect Ras-like protein mRNA or DNA.

Nucleic Acid Arrays

The present invention further provides arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee et al., PCT application W095/11995 (Chee et al.), Lockhart, D. J. et al. (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. et al. (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their

entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown et. al., US Patent No. 5,807,522.

The microarray is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides that cover the full-length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm that starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler et al.) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the

surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray so that the probe sequences hybridize to complementary oligonucleotides of the microarray. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of one or more of the proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention. Figure 3 provides information on SNPs that have been found in the gene encoding the Ras-like protein of the present invention. SNPs were identified at 20 different nucleotide positions. Some of these SNPs that are located outside the ORF and in introns may affect gene transcription.

Conditions for incubating a nucleic acid molecule with a test sample vary.

Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the human genome disclosed herein. Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid. Preferred kits will include chips that are capable of detecting the expression of 10 or more, 100 or more, or 500 or more, 1000 or more, or all of the genes expressed in Human.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers

allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified Ras-like protein genes of the present invention can be routinely identified using the sequence information disclosed herein can be readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in procaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory

control region to allow transcription of the nucleic acid molecules from the vector.

Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and

eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enteroRas-like protein. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene*

69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in a host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance, propagation, or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a

regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced, or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or

may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell- free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as kinases, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

Where the peptide is not secreted into the medium, which is typically the case with kinases, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a Ras-like protein polypeptide that can be further purified to produce desired amounts of Ras-like protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the Ras-like protein or Ras-like protein fragments. Thus, a recombinant host cell expressing a native Ras-like protein is useful for assaying compounds that stimulate or inhibit Ras-like protein function.

Host cells are also useful for identifying Ras-like protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant Ras-like protein (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native Ras-like protein.

Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a Ras-like protein and identifying and evaluating modulators of Ras-like protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the Ras-like protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the Ras-like protein to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse*

Embryo, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al. PNAS* 89:6232-6236 (1992). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al. Science* 251:1351-1355 (1991). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect ligand binding, Ras-like protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* Ras-like protein function, including ligand interaction, the effect of specific mutant Ras-like proteins on Ras-like protein function and ligand interaction, and the effect of chimeric Ras-like proteins. It is also possible to assess the effect of null mutations, which is mutations that substantially or completely eliminate one or more Ras-like protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention, which are obvious to those skilled in the field of molecular biology or related fields, are intended to be within the scope of the following claims.

SEQUENCE LISTING

<110> YE, Jane et al.

<120> ISOLATED HUMAN RAS-LIKE PROTEINS,
NUCLEIC ACID MOLECULES ENCODING THESE HUMAN RAS-LIKE
PROTEINS, AND USES THEREOF

<130> CL001188

<160> 5

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 3257

<212> DNA

<213> Human

<400> 1

tgcccgctgc	ccgcccgcag	ttcccggccc	cgctggcccc	agtcattggcg	aagcagtacg	60
atgtgctgtt	ccggtctgtg	ctgatcgggg	actccggggg	gggcaagacc	tgccctgctgt	120
gccgcttcac	cgacaacgag	ttccactcct	cgcacatctc	caccatcggt	gttgacttta	180
agatgaagac	catagaggta	gacggcatca	aagtgcggat	acagatctgg	gacactgcag	240
ggcaggagag	ataccagacc	atcacaagac	agtactatcg	gcgggcccag	gggatatttt	300
tggctctatga	cattagcagc	gagcgctctt	accagcacat	catgaagtgg	gtcagtgcag	360
tggatgagta	cgcaccagaa	ggcgtccaga	agatccttat	tgggaataag	gctgatgagg	420
agcagaaacg	gcaggtggga	agagagcaag	ggcagcagct	ggcgaaggag	tatggcatgg	480
acttctatga	aacaagtgcc	tgaccaaac	tcaacattaa	agagtcattc	acgcgtctga	540
cagagctggg	gctgcaggcc	cataggaagg	agctggaagg	cctccggatg	cgtgccagca	600
atgagttggc	actggcagag	ctggaggagg	aggagggcaa	acccgagggc	ccagcgaact	660
cttcgaaaac	ctgctgggtg	tgagtcctgt	gtggggcacc	ccacacgaca	cccctcttcc	720
ctcaggaggc	ccgtgggcag	acaggggagc	cggggctttg	ccctgctgct	gtcctctcgt	780
gtgatgaccc	tattgagtat	cagtagccac	tactccccct	gcctggccct	gagagcggct	840
ctgctgtcat	ctcaagcagc	ccctgtcccc	agcccgtcca	ccctggagtg	gtcttcttca	900
gcctgtttcc	ccagccacag	gcctgtctac	acccccacga	tgtgccgcaa	gcactgtctc	960
accatcccgc	acccaccaga	caacagccag	ggctggagtc	caggccactt	tcagctgctc	1020
ctttctccgt	gcacgtgtgc	tcttctctgc	tttttctctc	ttccccact	tctctttctc	1080
tgaccctcc	cctccggtgc	gtttcgtatc	aaagctcctc	aaaccccgtc	ccccgtgtgt	1140
cctgctgtgt	gcagctcgtc	ctttccttcc	ttcctaagct	atccaagggg	atggaccacg	1200
gctcgtgggg	aggttccacc	cttggatcca	ggaagaaccc	tccaccctgc	ctcgtgggtg	1260
ggccaaaggc	tacagggtgc	ttcttcctct	tccccacccc	ccactgtccc	tcagtgtcca	1320
tgggcctgcc	tccccagtg	cctgcgaaag	tggagcatcg	aggtaggagg	gaaacagcaa	1380
ccggggagtc	ctcgagcctg	gggctgccct	acctctaccc	attccccgac	cagagctttg	1440
cccttgcttg	gctgcccgcc	tgctcttttg	gggaactgag	ctcagaggca	ggtgcttcag	1500
agaaggaaac	aaaatgaggg	gtggcaggga	taaaaagtca	cctccattct	ctacctcca	1560
tgacagatga	acacaatttc	tctccacctg	gctcccaaag	ttaaagatgt	ggaccaaggc	1620
ctgtgggtac	tccaggggca	aggagagccc	tggggtcagt	gacactgtca	ggccaaccat	1680
gcactccaca	aaggggagca	tttggaatat	aaggactagc	tcctatgtat	cagggttaaga	1740
gcaaggagga	gctggccagg	gacagcagtt	tgacacagcag	aggggaatgt	agcaacagca	1800
gggcctccta	ggccccatct	tccatttctt	aggtaagaag	agcatttctc	cagactccca	1860
ggcggaggac	tgagcctagc	cttcagcaac	caagggtctc	ctgggaccca	aagtttatgg	1920
gagaagggca	aagacttcat	gggaagagag	aagggaaggc	ctgggtagaa	acgcttggtg	1980
ctgttctctt	tggcctttta	gacaaagcgc	tcactcttgc	ctctacctcc	tgataggctt	2040
gagggtttgc	caaccacact	gtggctacag	gtggagggaa	gaggactcct	tcctccagag	2100


```

tgctatgttc aggaagtttc tttaacccca tatggcccaa gagtagctcg taggaggccc 2160
tttaaagacg gaacaagtaa tttaccagtt ctactggggg tcttgccac cgtcccaagg 2220
tgggcgaggg ctaggaagag ggtcattctt aagccacaca ttagctgcac tgcgtggctg 2280
cagccaaaac aaagaactgg gtgttgagta ttcataact aagaaccaa atccagggca 2340
ctcatatgtg aaggataaga acctcacttc cttactcctc caaaaagaag tggggaaaga 2400
acctatcaac ctttcctcct gacttaccaa accaggaaaa cagcaggaga ggggtggctca 2460
ggacttaggg acagggtata gcttagatgg tggaaagcaa aggagagcag gaagttgtaa 2520
atcactggct aatgagaaaa ggagacagct aactctagga tgaagctgtg actaggctgg 2580
agttgcttcc ttgaagatgg gactccttgg gtatcaagac ctatgccaca tcacactggg 2640
gctagggaag taggtgatgc cagccctcaa gtctgtcttc agccagggac ttgagaagtt 2700
atattgggca gtggctccaa tctgtggacc agtatttcag ctttcctga agatcaggca 2760
gggtgccatt cattgtcttt ctctcctagc cccctcagga aagaaggact atatttgtac 2820
tgtaccctag gggttctgga agggaaaaca tggaatcagg attctataga ctgataggcc 2880
ctatccacaa gggccatgac tgggaaaagg tatgggagca gaaggagaat tgggatttta 2940
gggtgcagct acgtcacccc taaacttttg gtggcctggg gcatgtcttg aggcccagac 3000
tgtaagcag gctctgctgg cctgtttact cgtcaccacc tctgcacctg ctgtcttgag 3060
actccatcca gccccaggca cgccacctgc tcctgagcct ccactatctc cctgtgacgg 3120
gtgaacttcg tgtactgtgt ctcggtcca tatatgaatt gtgagcaggg ttcactctatt 3180
ttaaacacag atgtttacaa aataaagatt atttcaaacc accaaaaaaa aaaaaaaaaa 3240
aaaaaaaaa aaaaaaa 3257

```

<210> 2
 <211> 212
 <212> PRT
 <213> Human

<400> 2

Met	Ala	Lys	Gln	Tyr	Asp	Val	Leu	Phe	Arg	Leu	Leu	Leu	Ile	Gly	Asp
1				5				10						15	
Ser	Gly	Val	Gly	Lys	Thr	Cys	Leu	Leu	Cys	Arg	Phe	Thr	Asp	Asn	Glu
			20					25					30		
Phe	His	Ser	Ser	His	Ile	Ser	Thr	Ile	Gly	Val	Asp	Phe	Lys	Met	Lys
			35				40					45			
Thr	Ile	Glu	Val	Asp	Gly	Ile	Lys	Val	Arg	Ile	Gln	Ile	Trp	Asp	Thr
			50				55				60				
Ala	Gly	Gln	Glu	Arg	Tyr	Gln	Thr	Ile	Thr	Lys	Gln	Tyr	Tyr	Arg	Arg
65					70					75				80	
Ala	Gln	Gly	Ile	Phe	Leu	Val	Tyr	Asp	Ile	Ser	Ser	Glu	Arg	Ser	Tyr
			85					90					95		
Gln	His	Ile	Met	Lys	Trp	Val	Ser	Asp	Val	Asp	Glu	Tyr	Ala	Pro	Glu
			100					105					110		
Gly	Val	Gln	Lys	Ile	Leu	Ile	Gly	Asn	Lys	Ala	Asp	Glu	Glu	Gln	Lys
			115				120					125			
Arg	Gln	Val	Gly	Arg	Glu	Gln	Gly	Gln	Gln	Leu	Ala	Lys	Glu	Tyr	Gly
			130				135				140				
Met	Asp	Phe	Tyr	Glu	Thr	Ser	Ala	Cys	Thr	Asn	Leu	Asn	Ile	Lys	Glu
145					150					155				160	
Ser	Phe	Thr	Arg	Leu	Thr	Glu	Leu	Val	Leu	Gln	Ala	His	Arg	Lys	Glu
			165					170					175		
Leu	Glu	Gly	Leu	Arg	Met	Arg	Ala	Ser	Asn	Glu	Leu	Ala	Leu	Ala	Glu
			180				185					190			
Leu	Glu	Glu	Glu	Glu	Gly	Lys	Pro	Glu	Gly	Pro	Ala	Asn	Ser	Ser	Lys
			195				200					205			
Thr	Cys	Trp	Cys												
			210												

<210> 3
 <211> 28770
 <212> DNA
 <213> Human

<400> 3
 gctcaagatt gcacagctgg tgagtgggtga cactgggact ggaacccaag tgtgccttac 60
 tccagagccc ttggcatgca cctgaaaccc catgtaagcc cactgtggag acgcgcacct 120
 cgaaataatg gaatccaacta catcagttcc tttagctttc tgtgtaatca gagtagctag 180
 caggctcggg atttcgcccc cgggcttttt tttttttttt tttttgagac agagttttgc 240
 tcttggtgcc caggctggag tgcaatggcg caatctcggc tcaccgcaac cttcgcctct 300
 caggttcaag caattctcct gcctcagcct cccgagtagc tgggattaca ggcaccggcc 360
 accacgcccc gctaattttt ttatatattt agtagagatg gggtttcacc atgttggcca 420
 ggctggtctt gaacttttcc cctcttatta taattcagac acttaacctg aaatatacct 480
 tttcaaataga agtaaattggg cttaccactt tccttgacct actattgaaa aatacattct 540
 ccatccaata ttcagcctga aaacagggtat gtacatatat acttttcatt gctttttttt 600
 tttttttttt gagacaagggt ctccctctgt tgcgcaggct ggagtgcagt gtcattgatct 660
 cggctcactg cagccttccc ctaatgggtt caagcaatcc tcccacctca gcctctcaag 720
 cctgggatta caggcgagcc accgtgcccc gctaattttt ttttattttt agtagagact 780
 gggtttcaact acattggcca ggctgggtct cagctcctga cctcaaagtg atctgcccgc 840
 ctacgcctcc caaagtactg ggattacagg catgagccaa cgcgcctagc ctttcattgc 900
 tttttaaaga cctaataaggc tagactttgc tctccctcaa tactcgttgg tagggatagg 960
 caattttctc aactccggag agcattcatt tgcctctctc cgggtgctaac acattcagt 1020
 gtaggaaact ggatcttgaa caagggccat tcatcttttg gtgccactgg ctataccaca 1080
 gagaaattta ggggtctgaa acaatacatt ggtcacctgg gcaacctatc taagcacctt 1140
 agagggaaaa cggagacttg cccgcacacc tctaaaggat tttgcaactg gagatgttct 1200
 tatggccatc tatcttttca ccctggtgga ggccgtgaat aggcattttc cccatttaaa 1260
 gaaaaaatgg ggacggggga gggccgtgac acagtcacac aggtaagggg cagccagatg 1320
 gcaggagggg ggaattccac ccacactctc ggggactcat ggagacgggt gttcgaatcc 1380
 agatcctgct caaggccttc ctacctcggg tgagcccagc tgaggtagca gccactgggg 1440
 agcccggcca gatcctgcag atgcagggtg ccacggcggg cgggaattacc ggcgcagac 1500
 ttgggggtgg atatggggag aagtgtgtag cccggaaaagc ggagcacggg agaagtgggc 1560
 tgggtggggg ctcacctcaa ctccccatt cggagcgtcc gcggaaaaac gaaaacgttc 1620
 ccccgccccg ggcaggaagg ggttgggagg gggggctggc gccccgcccc agcgtcgctt 1680
 gctcgatggg gtcccgctct cctgcgcgcg ctccccgccc cctctctacc ggggcggcgg 1740
 cggcggcgca ggggaagggg cgggcagggg ccgcccggcg tttctcctcc caccgcctcg 1800
 cgccagcccc gccgagccga gccgagccga gcgggcggcg cgccgggctc ccgcccagc 1860
 cgcgcttccc ggcacccagc gagcagtggt gcaggcgggc gggcgaggca gccgcggggg 1920
 ccggggcccg cgtctctctc gccgcccga gcgtccccgg gcgggcgcgg gccgcgatgg 1980
 cagcggcgga gcagggtga gcccgctgcc cgcgcgagc tcccggcccc gctggcccca 2040
 gtcatggcga agcagtagca tgtgctgttc cggctgctgc tgatcgggga ctccgggggtg 2100
 ggcaagacct gccgtgtgtg ccgcttcacc gacaacgagt tccactctc gcacatctcc 2160
 accatcggtg aggggcgggt gcccgggcg cccctcctc cccgcccgcg gcccttttcc 2220
 ccgcccggcg cgtccccagc tggggaggaa ttgccagccc ctccggctgg aggcgggtggc 2280
 gccggaggcc ggagtccggg ataaatctcg ggggtgagcat aggttttggc aggtgagggt 2340
 gtccctgctg cctgccgtcc ggaccagggg tggggctctc cgcctcttgc cgggaagcct 2400
 tccgtcccat caaacgaga aaccgggggt gaggggagct ggtgtaggcc tgggtacccc 2460
 gagctggggg agcaagaatc gtagccgctg gaataacacc cccacacccc cagggggagg 2520
 ggaagtaaag cttctgtctac ggaaaagggg gtcagggtgg agaccggagt cactgaggcg 2580
 cccttggttc tgtggtgacc caaggtggag ccggcggggg gcgagggggg gaagagaggga 2640
 cgtacggagg ggcacaggg atcgagtttc cagggcagag ttgggaagggt aagccgcaag 2700
 gtgggacacc tgggggagga cacagatagg gtgaggagcc cctgcgcctg ggaagaggag 2760
 acatctgttc tgaggagggc taaagaggat ggaggagtgt caggaatacc tgcccagacc 2820
 aaggggtcag aaggcaggca ggaccgcctt gagggcatct ctcatctggc agtgctggag 2880
 cctgtggtta gagggacaag acccggtggc atcccagaca gcactatgat ggggtcactt 2940
 attctaggaa tgggtccatg gcctccctc tgagacagtc agtctcccgc ttctaggctg 3000
 tgagggggccc cctccctgag aagtctgagt agagggaatt tcatcctcag ctgctacccg 3060

ggtcagccct ggagtagcct ctgcattgcc caagcccctg gaaacacctg ctggctggct 3120
 ggtcatccat ttggaatgct ctccatagaag tccctgctgc catcagggat gggcaccagc 3180
 tctcagcttc ctcttgagga ttcatgtcca caccatcccc cctcccccca acacacattc 3240
 cttgctgaga gagaagtagg agcagataga tacagccagg aggaacagaa ccttctgggt 3300
 aagaagccag ctttattgtc caagagacct gagacctcac tgtggggcaa agcaaccttg 3360
 aatattgcct aaacttctga gctttattta gtttctcatc tgtagaacgg gtataataat 3420
 tgcacctacc tgccaagtgt ttgtcaagat taaatgagat aacgattgtt aagtgccttag 3480
 cacagccaga cacatgggtg agctcgataa atgctgattg ttcttactgc tattgccatt 3540
 atcattgagc ttttaggggt tctctctctt gtttcaccaa cttgaagggt gaaacaacag 3600
 gacttagggg cagggaaacag aacttgtccg tctttctcag aggagctgta aggccaaactc 3660
 ttaggaaacc caggagcttg ggctgagcca tggtttgat gagagacatt gcagaaagaa 3720
 ggggagccta tagacactaa ggctttgtgc ctgccgggag gacttgggga agaggcagggt 3780
 gcaggagaaa ggcattggcg tgatggagga agtggcagag gaaccagatg gtgtatgagg 3840
 acaggttgtg ggctcaggga caaagggcgg tgggttatcc ctttaaggaaa ctaggagtgg 3900
 ctatttttgg gagaggcctg gtgcttgga ctactgagct atctccagag agctgtgggc 3960
 tgcctgggag gccctggctt tggcctgaag agctgttgtt tgcacctgct ctccatagtc 4020
 cattccaagt cctatagggt acatggactt ttccctttga gggcttcatt caaccacctc 4080
 atttcagaag ctctgggact cctgcttagt gctgtgggag gcagcctccc ctgggagaca 4140
 cataccctcc tttttgaggg caccctctt tctaaaatac caggatggcc ctctgaggct 4200
 cgtgctctcc ttaaagagag tccattgcct cacacctcta atcatccacc cttctccttg 4260
 tcccttcccc ttgtaatctc ccttcttaga cacttctgc taatagggtga aactaaata 4320
 ggtcacaggg acttcctgaa accctccagg gcagaccact ttgggcacat aggtgaatca 4380
 gtgaactgag taggggtgtc tctgcagcac tgtctccct caaggccctt ggtatattgg 4440
 cctaaaacct aaagatggct ccagatttct tctctccgt tctgacaccc gggttccct 4500
 ttctacagga cacagaggat tagaaagaaa cctgtcttct ttcacaccag ctttttaaaa 4560
 ggagtttgga ttccatggaa tctctagggt ccccttttcc acaggacaca gaggactcta 4620
 tctcccttat cttgggcgtg tttttcagca ttaactaaaa cttaaaggaa agagttggat 4680
 ggtcaagaaa agctttttcc ttaagtata tggacagttt ctcaaggagg tagaaggggc 4740
 agccaggaga caaatcaagg agccaacgaa atgagtgtca ccaagtcata gtcattcgct 4800
 tatttttaaa aaatgcgtgt cctgtatgcc aggtctctga ctgagaccga gagattccaa 4860
 gatgaataat acctacagtc actgttctca aattgtgcat tacctaaaac acattacatg 4920
 accatgctgg ccactgatcg aggcacctt cccaggggct ttttttgtga attaagaaaa 5040
 caaggttaatt caccagtatt tgccaagata gtttggttcc ttggtcatg tggatatcac 5100
 ctaggccagt acttttgtga tttactgtgt actccacttt aacggcctgc gatcttctag 5160
 agaagaacct gccaggaggc agtgagaggc ctccctggta gactgagaca ctgactgtcc 5220
 ctccccctat ccttttcgtc tttctggcca gcagaccagc aggtggccct gccactggct 5280
 ctgccacagg catttcttct ctgtgcagct gtgctggcct ggctgggggt tgggtcggaag 5340
 ggggtcccaa aatactacct taaacaaatt aattgagcat tccactacca gctctgtgcc 5400
 aggcatttta gagacatatt gcagtctacg ttttctgccc acagaagccc ataacctaga 5460
 tggggaggga agacaaaggg aaaaaacaaa aacaaagagc tagtgccaaa atgagatatc 5520
 tgaaagaact tggatgaatca ctcttcaaat gtaaaggatg gattatgatc attgcagtta 5580
 ctcttaatag aggtctcaca gtgggtatca gaagctaaat tatgatgcaa gatgtaccat 5640
 gaggcagccg gagaatggcg atggatggga tgggtgagtg ctattcccac gactccatgc 5700
 tgtcggaggc tggggaagag agaggccct gtggactaga accggcaggg aaggctgaag 5760
 ctaggcctca gtgtgggctg ctctcagtt cctgcagcag aaggagcag atggagtaac 5820
 atgagcagag ataacagagg tgggattgag taggtgtccg tggggtctta ggcagtttag 5880
 atgcaacaga agggattctt caggaaagtg agaagattct tctgtttctc tctctgtctc 5940
 ccaaattata agtgccttga tgggtgcgacc aaatcttatt cctcattgtt tttatagtc 6000
 ctagtacagg gccaggcaga ttcaatgcct gttgttaaata taatgaatga atgcagggac 6060
 cagttggcag agggcattga gagcctggcc aaggaggtgg aacatgagcc ttagcaatgg 6120
 taggaggggt tttgagtagg gtactaatga ggttggctgg aagaaggggt taagacttga 6180
 agcagggaga ctagtccagg gctgcagtag tctctgggc atgaaggaa ctctgaatgg 6240
 cccctcacc ccagtggtag caacaccaac ttccacacag tcagttgttc tactttccct 6300
 ccagaatggg gagtgggtca agccaatcaa cctggcaact tctgaaagaa tcttatggga 6360
 cctgtgccat gaccaggtag ggagaagatg tcatacatgg acatctatgt tcaggggacc 6420
 tttgaggacc tttctgcatg gtggccagga gtgagatgat gtaaaccaca aatggaaact 6480

gaagagactg ctcaggagtt gttgggttttc ttttcttttc tattttttttt tttttgagac 6540
 taggtttcac tctgtcacc ctcaggagtt gttgggttttc ttttcttttc tttttgagac 6600
 gcctcgatct cctaaacgca atcctccac ctcaggctct caagtagctg ggactacagg 6660
 tgcattgccac cacattcagc taatgtttgt acgttttgta gagatggggt ttactatgt 6720
 tgaccaggct ggtctcgaac tcctggactc gtgatccacc agcctcagcc ttccaaaatg 6780
 ctgggattat aggcgtgagc tacctcactc cctcaggagt tggttttctc cctcccatcc 6840
 ttagtcttcc ctgagtagac ctgtcaccta gtccctggac cttttgtttt gaaagccacc 6900
 ctccaggcta cactccttct ggggtgaggag gagggtgatc tggttggaca ggttgggctg 6960
 ctgtggcttc agggcacttt ctcaggctgg gttgctgctg ctatgtcacc tttctcaagg 7020
 agttctgctg ggactggctt ggctgcctgt cttgactttg cttttgactg aggagtgagg 7080
 agatgggtgag ggaggggggtg gggctagatc caagcctgga atgggggtgac ctaacagaca 7140
 ctggggcctg tgcttagaca ctaggatcct ggggtttgca ggtttctaga ctgagaggag 7200
 ctgggggcaa atgcagtgtg acgtttgtgag agggctcagg ctgggtctgt gtcagccttc 7260
 aggcagcctg agaccagtct ctacctactc tgttcccctg gtacctagaa aggaagggaa 7320
 ggtgagaagc aatgagcaga atggaaagag cccagattaa catgcacatt tcccatggcc 7380
 ttactggccc tgtgaccttc agacactttg atgacatctt tgtgcttcgt ttctgcatct 7440
 gtaaattgaa gatggttaaca gagtctttct taaaggttgt tgtgaagatt atagagccta 7500
 gcgcataata agcacttggtc agagccctcg ataaaaataat agctgctatc atattatcat 7560
 tattattatt ttattttatt atttatttat ttttttttga gaccgagtat ctctctgtcg 7620
 cccaggctgg agtgacgtgg cacaatctcg gctcactgca acctccatct cccgggttta 7680
 agtgattctc ctgcctcagc ctccctaagta gctgggatta caggcaccca ccaccacacc 7740
 cggtatttat tattattcct agctataaga atgctgtaga gatgaataca ctgtcagtga 7800
 gctaggaggt catcctgtgt atccatcact tgtgactca taagatttgg ggacacagtg gtgacaaaa 7860
 tgaacaccaa ctacatgcca ggtgccatgc ccaagggagt gcataggaag ctgtgtttca 7920
 cagacagaaa ccaaggagct ggcttacatt ccaagggagt gcataggaag ctgtgtttca 7980
 tttcagtttc tgctctagta ccccccttcc cctggcagtg ccagggtctg agaaggaaga 8040
 gtgaggtggt gagggaggtgt gaagcagtgg ggtgacctga gaggagagga tgggggtggct 8100
 ttgctcaag gcttgggccc ctgctagggtg tctctctgcc tcaggcctct gtttctcctc 8160
 ctgacacagg cacagactcg gcctccacc ccttcccaaa ggacatgacc ttgggaagga 8220
 acatatctga agcccgcgga gggtttccgc tgcctgtgcat ctgtgccaca gatccgcaga 8280
 tgcaccaca gctgggagca ccggttcttc ccgcctacct gcactccctg gtttctgttc 8340
 ctctctctc ctcttctctt ctccccgctc cccagacagg ctggtgatga gctttataac 8400
 atgaaagctg atatttggcc attatccttc taccctgatt gccagctctt ctcagagtgc 8460
 ctctctctgt aatccaatct ttgcaccagt ttccctgtga aactgccagt tttctgtata 8520
 ggctctgccc ctctccttgg ctcttctctc tggtcagtga gctttgtcaa ggggaacaca 8580
 gggcttctct gacacgtaat tcctccact gaggaggaag gggctaataa ccagccctgt 8640
 tttattttat tttatttttt tgagatgaag tctagctctg tcgcccaggc tggagtgcaa 8700
 atggctcgat ctgggtcac tgcaactctt gtctcccggg ttcaagcgat tctctgccc 8760
 cagcctctct agtagctggg gattacaagc atgcaccacc acacctgggt aattttttgt 8820
 gtttttagta gagatggggg ttcaccatgt tggccaggct ggtctcgaac ttctgacctc 8880
 agctgatcca cccacctcgg cctcccaaa gctgggatt acaggagtga gccaccatgg 8940
 ctggccgacc catctctta aaaaaacaaa aagaaaaaga aagaaaaaca aacaaaaaca 9000
 ttttttaaat taactgatta tgggtggcatg tgctgtagt cctaactact caggaggctg 9060
 aagtgggaag attgcttgag cccaagtagt tggaggccac agtgagctgt gatcacacca 9120
 ctgtactcca gcctgggtga cagagtgaaga cctgtctca ggaaaaaaa aaaattactg 9180
 agaactctgt gacctgggca ccatgaacta tagaaagggc taacagttgg ctttgaaatg 9240
 tgggttatgg ctgggtgagg tggctcacgc ctgtaatccc agcactttgg gaggccaagg 9300
 tgggcagatc acaaggtcag gagtttgaga ccagcccggc caacatagtg aaacctcatc 9360
 tctactaaaa atacaaaaaa ttagccgggt gttgtggcag gtgcctgtaa tcctagctac 9420
 tcgggaggct gaggcaggag aattgcttga acccaggagg tggaggttgc cacaagctga 9480
 gatcgacca ctgcactcca gcctgggcga cagagcaaga ctccatctca aaaacaaaaa 9540
 taaaaacaaa aaaaagtggg ttgttttctt ttctttcttt tttttttttt 9600
 ttttgaaaca gagtcttgct ctgtcaccag gctggattgc agtgaggat ctgagcacac 9660
 tgccaacctc gcctcccagg ttcaagtgat ttccctgcct cagcctccag agtagctggg 9720
 actacaggca cgcaccacca cgctgggcta agtttttgta ttttagtaca gaaggggttt 9780
 caccatgttg gccaggatgg tctccatctc cctgacctcg tgatccgccc acctcggcct 9840
 cccaaagtgc tgggattacg ggcattgagc accacgcccg gcctaaaagt gggttatttt 9900

ctaattgctc ttccctgatt aaaattttct ctttgcccat cttttctcta gatatgtact 9960
 gacttcattc atccatttat tctgtctcact tgctcattca tttttgcttt catttactct 10020
 actttgttga ataataatata gtgatctacc tgctgccagg cagttagagt ctgaagtga 10080
 caggatgctg ctttgccctc tgggagctta cagttagagt gggaaccaga catccaaaca 10140
 agcagaatat tatgcaaaaag aaatgtcagg atgctttgga atcacagagg agtgagaaat 10200
 ccctcccggg gaggtggtg aaggctttga agaggaagtg acatttgagt ggagtcttga 10260
 agactaggca ggattctcca ggggccctgg gtgtggggga agcacacatc ctctccctg 10320
 taggaggtgc tgtggagaac acctccagtg gggctgctac tcttcagcct tgctggggcc 10380
 agctggagtg gccacaccat ggtcacacca gctgaagtgc aagaagcccc ttgccaggag 10440
 attgctttgc tggctctggg tgagggcagg tgcattctga agcccccttc tttctaagat 10500
 gtttgcctct gaggttctat gtcctagtct tttcttccct gaaccttttg ctaccagtca 10560
 gcacagccct gcctgagaag gaggtctggg gagttagtgg tcagttagcct ggtgggtctt 10620
 ggctgcctct gtggtgcccg ctggcctaag tagcaggctt agggaggcga gaccagttc 10680
 caggggctgc caatggggag cgagatgggg tggctggagc aactgcaca tgtcaccaag 10740
 gctctaggga ggtctgtgca caaggcagtg ggaagaaagcaa ggggaagacc cagcctggtc 10800
 aacatggtga aaccccgctc ctactaaaaa taaaaaatt agctgggtgt ggttagagcac 10860
 gcctgtagtc ccagctaact tgggagcctg aggcaggaga atcactttaa cacaggaggt 10920
 ggaggttgca gtgagccgag atcgtaccac tgtactccag cctgggtgac agagttagac 10980
 cctgtctcaa aaaaaaaaaa aaaaaaaaaa aaaaagtggg gaaggggaac actgactctg 11040
 attatctact ccatatactt actatgtacc tactacctac acagggacgg tgggctttac 11100
 gcatgccatt tattcagtgt atagagatct cagcatcaca taggagcagg gaggttctga 11160
 gttggccttg ctggcatttg agaagtttct tgggtgattc ttcaggttca cgcctccaga 11220
 caagtgaag tgctattgaa tgctgactat gttccaggaa ctaaacacaga tgctagaaga 11280
 cacgcagtaa acagtacaga tgcagggtga catgtgaggg tccacacaag acctgagaga 11340
 agggaggggt ctgtgtcag tccccctttt gtaacaaagg agagagtact gttgacctc 11400
 ttcttaggaa ctgtgagttt gactgaaatg tgtcctgcca caggatcttt gctgcttct 11460
 ctacctgatt ctttgagctc ccctgctggc accttcttgt catttaggtc tcagctcaaa 11520
 tgttacctcc tttaaaatgt cttctctggc cagccagtct aagggtgctt gtgcttgggg 11580
 tctctcact ctctacttta tccgcagtt gcttcttctc acatatggct ctctgaaatt 11640
 aggtattcat tacttacatc tgtcttcccc actagaatta agctctgatg acaaggatct 11700
 ttctgtgctg ttcatagctt atcttctagt acctggctta gttcctggca cattgtaagc 11760
 attcaataac agtttgaatg aatgaattaa caaatgaagg aatgaatgaa tgcattttcc 11820
 tagaggactt ctgttcttcc ctgagggaaag ttataggctg tattgggtttc ttgggactgt 11880
 tttttgtttg tttgttttgt tttgtttttt gagacagagt ctactgtat cccccaggct 11940
 ggagtgcagt ggcacaatct tggctcactg caacttccgc ctcccagggt caagcgattc 12000
 tcatgcctca gcctcccag tagctgggga ttccaggagc ctgccaccac gaccagctaa 12060
 tttttgtatt ttttagtagag acaaggtttc acctggttg ccaggctggg cttagaactcc 12120
 tgacctcagg tgacctgctt gcctctgctt cccaaagtgc tgggattaca ggcatgagcc 12180
 accacgcccg gcctgttttt tttttttttt taagacagag tcttgactg tctcccagac 12240
 tggagtgcag tgggtgtgat tcagctcatt cgagctcaa cctcctggcc tcagggtccag 12300
 gtgatectct taactcagtc ttctgagtaa ctgggcccac tggatatata caccacacct 12360
 ggctaatttt taaatttttt gcagagacat ggtctcacta tgttgccctg actgatcttg 12420
 aactccttg gttcaagtga tctcacacc ttggcttccc aaagtgtctg gtttacaggt 12480
 gtgagccacc atgcctgggc ttgagactgt taagatgatg aggtggagg gagtggatgg 12540
 cctcactgct tgagccctag agattcctta ctccagagt cctggctgc agaggtggcc 12600
 ctggagggtc actccagcaa cctggctgag ctgatgggca tcatctgata ccagctctga 12660
 ccctgaataa taggcaacat ggaccttagt ctaaagcact gacctctcat ctctgcata 12720
 accaaagaag atgagatttg ggtgaggaca cagccaaacc atatcagctc ccgggatccc 12780
 tgtgtgaatg gggctctttt tgggtgttga gggctgcaca gggtgacctc tttagagggt 12840
 acctcctgcc acaaccaca ggaggtgcac atggcccaca catgctgggt tctgacgtg 12900
 ggaggggctg gggcactcct gggacctgtg cttggtaact ggagctggcc tggccctggg 12960
 gattgggtgt ctgccttggg tttcagggtg attaggttgt tctctgttgt ggagtctcat 13020
 tactaatgaa aagttcaggt cgcactgctg gtcctttggg ctgtggttga tctgggtgat 13080
 aacatttggc acccagaggc agccctgttt ccactgaagc atgcggagct tggctggcag 13140
 gcaggcaagc tggcagctgc ccttaaccca tgaggtgctg gcccgtagt aggcacacc 13200
 tacctgtgcc agaattgagg ttgtagccag actccaggag ccactctggg cccacagggg 13260
 gcggcatttc ctctttttgt tgaaacattc cagccaagtg ctggcttggg cttcatctct 13320

ctgtccact ctccttcctc tccccaacat aagcctcctt ctacatccta gagctctttc 13380
 cattccccct cctgcagctc tgggctcgct aatctcatgc ttccctaagg gagcttgacg 13440
 gctgcttctg ctaacattta ataaagtctc gcgtgccaga ccctgtgtta tgggttttac 13500
 accttatctc acaatcttaa aaaaaaaatt ctctgagaat cctctgtcac cccacttta 13560
 caggtgagga aactgaggca aagataggct aactggcttc cccaacacca tgcaggtaat 13620
 tagtgataaa ggcaggggtg gaaccaaact tgacctccca attgtgctct taatggccag 13680
 gacactctgt gtcttgagcc acacttcctc catgttttct agggctttct agggaggcag 13740
 acagtgatgg gaaggggtgt tcttttagtg ggatgtgccc tgctgtctcc tttctgtaag 13800
 cgtcacagca cctccactgc tgtactgggg aggcaccaag ttttccctg tttgcccacc 13860
 caaggcgagc tagcttagga gtcacgtgag tgctgggtgt ctgcctgct gcatccctct 13920
 atcctgcccc tgcccccggt gccagagga gggccctgcc tgtcttccca gttctccaac 13980
 agcagcgctg tcccagcacc ctccgggctcc agttgtggcc tggcagctgc tggggcagac 14040
 accatacaga cagagtcaca gcaggaagag gatggggccc agggctgctg cctcaggcca 14100
 tggctgcatg gcaccatcag ttgattgagg agcttttctt gccaatgtct gaggcatcag 14160
 gtggcaggac acgtctccct gctcttaagc ctcaggcatg cagcccttct tatgctctct 14220
 ggggtgaggg ggagatcccc ctcatggaat tgcttttttt tttttttttt tttttttgag 14280
 acagggctct gctctgtcac tcaggctgga gtgcagcctc aacctcccag actcaagtga 14340
 tcctcctgcc tcagcctccc gagtagctgg gaccacaggt ggacaccatc acacctgggt 14400
 ttttttgttt tttgtttttt gttttctaga gatggggctc cactttcttg ctcagtctgg 14460
 tctcgaaact ctgggcgcaa gcagtcctcc cactcgtct tcccaaagt tttggattac 14520
 aggtgtgagc cactgtgctt ggccttttta tttatttaga atttgttttg gaattgcttc 14580
 tttatgcctg gcaactatgt ggcactatgt ggcagagatt ttaaaaacga gcaaacaaaa 14640
 caaatgcttt gtcaaccaca aaatgtattc tctgcccctt aggttctttt tgtgtagttg 14700
 aggtagaag acaaaaatag ggggcagtaa ggagcagggg gcgatgggtt aggaggtctt 14760
 ccttcagcc ccttggttga agcatctggc tcactagctt gggggagcca ttaggcagca 14820
 gtggccaatc ctgaggcact ctcagggtgc actaagaaaa ggggcatgtg ctctatggat 14880
 acccatgggc tgaacttgga gtctggtctg gacctatggc tgtgctagga tccaccgtcc 14940
 ccagcccaa ctgcagtcag catgttcac atccttaggc ctctccgctt ctttctgcat 15000
 gttgtctgc ctcatgccct gctcattacc aactggctcag tccccactgc cctgcctgga 15060
 gtgagctggt ttgattggct tagctaagct ccttgccctc tgctggccag gtcaccctgt 15120
 gggtcaccag caaacctgtt gatggctcag tctgaacctg cttctccaca aagaacatgt 15180
 tgcaccagc cctgcttctc tgagcagagg tttggggctg agctgttcta gccagaaaagg 15240
 gacacaggg gtggcaggca ccatgatggg catatcta atctgtctga aaacaatgag 15300
 ctgctctccg tgctttgggc acctggttgg gagagggccc atctgtctga ctttctcctc 15360
 ctggggctct cagcgtctcc gagaacctct gccagagctg tgtagaagtg gtttgcttgt 15420
 ttcttaacac ttctgtgccc tatttctttc tgtacccaag aaaggaagta gactgttttg 15480
 tagggacact gtcgggggtga tgaatctgga cttactggaa tcatgaacca tgccaaggag 15540
 gaaggagaaa ataggctatg gtgggtgtct tagttagggc tggctgctgt aacaaaaatgc 15600
 ctttagctga gtaatttaaa gcaagagaaa tgtattgctc agagtttggg aggcctgggaa 15660
 gtccaagatc aggggtgccag cagattcagt gtctggtgaa ggctgatgct ctgtgacaaa 15720
 ggtggcacct tctagctcca tcctcacatg gcagaagagg gaacaagctc cctcagacct 15780
 cttttctaag ggcgttagtc ccatgcatga gggctcta acacagactg agtcacctcc 15840
 caaagccctc acctcccacc agcactgcac tggggattaa gtttcaatat gggaattttg 15900
 gaggaacaca gaccttcaga ccacagcagc gggcttctcc tcatgtgccc cctgcctcac 15960
 ttctagatgc cgcataatgt cagtgaacc cctctcttac taaaaataca aaaaattagc 16020
 tgggtgtggt ggcacgtgcc tghtaatcca gctacttggg aggctgaggc aggagaatcg 16080
 cttgaaccca ggaggcagag gttgcagtga cctgagatcg tgccactgca ctccagcctg 16140
 ggcgacagag gaagactccg tcaagaaaaa agagaaaagg catcaggtat gccaggggtg 16200
 gcgggaaaag gcatcgggta tgccagggcg tgtgggaaaa ggcatcgggt atgccagggt 16260
 gtgtgggaaa aggcacgggg tatgccaggg catgtgggaa aagggtggtaa gattcctcag 16320
 cctcccaggg ttgggaagcc tctggccgag tgaagcatac cctgggtggg ttttaagaca 16380
 ccagctttcc agtcagctc agctgtggga tgtgggaaca tgagtcagtg ggaacatgag 16440
 aattggcttc cctgtggctc acaataatac ctactcctgc ctacttcag ggaccgcat 16500
 aagagctgag ggattccata gctcaggggt atgctgtaaa gacaagcact atgcacctgg 16560
 gtgtgggtct gaaactttct tgtgcagaag agtgagtagg gctgggcgag tcctgagaat 16620
 gtgcatttct cacacacctc tgatgctgct gatgctctag tcccttggct ggcaagggtg 16680
 cctggttagt aggggccagg actctgtaat gccttccact tcagggttct ctgggctgggt 16740

```

tttcttgact cccaggaag cctttattca gcagagggaa ggtaggagtg agaggactac 16800
gctgtcagtg cttcacatac atcgtttaat ttatcccagc acagccctta ggaggggaagc 16860
agtattctcc ttctacactt aagaaaatcg gcctgggtcg gaggtcatg cctataatcc 16920
cagcactgtg ggaagctgag gcgggaggat cgctggagcc caggagtcca agactagtct 16980
aggcaatata gggagacctc atctctacaa aaaaaaaaaa aattagctgg gcatgggtgg 17040
gcacacttgc agtcccagct acctaccagc aggtgagct gggaggattg cttgagtcct 17100
ggaggatcga ggctgcagtg agctatgatt gctccactac actccatccc tggcaacaga 17160
gtgagactcc atcccaaaaa aaaaaaaaaa ttgaagctag gagaagtga gacttgccctg 17220
aagttacaca gtaagtgcc aaccaggac ttggaccagg tctttctgac tccaggccaa 17280
tggatgtttc ttccatgaca tatatagctc ttgaaactac ttctatctaa tatcaccac 17340
agtgtgttta aaaatacaga tttctgggcc tcacctcaa attatgattc agtaggtcta 17400
ggcagctcaa ggtcattgtt tttgtctttg ttttaagtca ccccagggtg ttctaaagcc 17460
gaagctctgc aaagcacacc ttgagaaaca gagaactctt gtgctctcgc tctcttgaca 17520
cttcagggtg aaaacttttg tcctaattgtc gttctcaaac ttacgcatgt gtgagaatca 17580
ctgtgagagc ttattgaaac tgattgcggg accccatacc tagagggcct gattctatag 17640
gtctgaggta aggcccaaga atttgcatat ttgcatttcg ttttcttttc ctttcttttc 17700
tttttttttt tttttgagat gaagtctcac cctgtcgccc agactggagt gcagtggcat 17760
gatctcagct cactgcagcc tctgcctcct gggttaaagc gattctcccc acaccccaga 17820
cccgtcctcg agtagctggg attacagggtg cccgccacca tgactagcta acgtttgtat 17880
tttttagtaga gacgggggtt tcacctggtt ggccaggctg gtctcaaact cctgacctca 17940
ggtgatccac tcacctcagc ctcccaaggc cttgggatta ctggtgtgag ccaccgcgtg 18000
cggccagaat ttgcatttct aacaagtccc aggtgatgct gatgctgtgg gtccagggac 18060
acactttgag aacagcttgt tactcaggcg atatgtggac agtagcgtca tcttcacctg 18120
ggagcttcct gcagcatctc aggccttgcc ctacacctac cagatcagaa tctgcatttt 18180
aactcaatcc ccgctgatt ctcatgcacc tggaaagttg agaaatatga ccttagagga 18240
gccggaatgt gaaaccactg gaggcagaga tagatggaga atatctcttc ttctcacgga 18300
tactaaagat gcaacaaaaa gggctgactc tctgggtgtg caccaggtg gggctgatga 18360
ccgaaaagag gccagatgtg gcagaggac tcttccctga gggaaaggcag agagaactta 18420
ggaaaatctg aagaaaggag gtggcttcag aggaaaggca ttcactctggg ccataaaaca 18480
gtggagaagg tatcctgctg agagcacagg ggtggggagg ggggtgccctg gagctgaagt 18540
cttcagtggg gggacagtga taggtgaaca cacatgtgaa taaacagttt gctaagcagc 18600
tgcgagggct ggccaagggt agaaagcatc cgtctgcaga ggcctcaata aggccagtgt 18660
gttgactttg tcctgcagtg ctgagcagtg gaaaaaacca acagccacgc agggagaggg 18720
aaggagccac gatgggcacg ggttactggg gccagggctt gactggtagg tggacacagc 18780
tgaaggccca ggttgtgtgg gaacagagcg cagaagcaat agattcctct tgaagatcct 18840
tgggctgtta acctttttta aatttaagag aggttgtgtg ggcggggagg gaggaaggaa 18900
aatccttcag aagacataga cttactctgt ttcttccatc atatgtgaat gcatatgaat 18960
agccaaaagg tgaataaaac acatgttccc aggtggccag tgagacctag gttgcaagat 19020
ggtgggggtg gtgtgaggcc ggggagtgct gcgagccccg gaattcctca gccttagtcc 19080
cccgccacat agctaagaag tgaggaggga ggtgagaagg agtcaactgc cagcctcact 19140
tccggtggag taccctgtct cctgtcagc tctgtctctg gggacagttg cctgctttca 19200
cctctccctc taacccctct tctctcacag ggaaaaattc acctaatat tggaaagttcc 19260
tctcctagca aagtccttct caggcaccca caggcaaaaa ggaaactaag cagagttagg 19320
gcttccaggc ctagccaact acacgactct cctcttgctt ccctaagaac cagcgcaagg 19380
ggcagcgtgg gttccagcat agatggacct gtgttggaat ctctgcacgt gctgtgctga 19440
ccctggctag ccattgacct ctctgagccc ttgtttcctt tccactaggc tctctgaggg 19500
caggggccat gtctttttca ctgctctgtc tgactgagc actgtgcagg gcacatagga 19560
agttcccata aatgtttgtg ggataaagga aataaaacct tctctcttcc tgtccccctt 19620
gtgatggctt tgcacaaggc actgtccttg gccaggtttg ctaggctagt gtgaggataa 19680
accaggtata ttacaaattg gagaaaattt ctcgttcttc ttggaagaag gtgctgtatc 19740
atgaaacaag aatgtcttga ttcccttcta tgccaggtag tggggagaaa cagggtgcctg 19800
ataaccgttg atccaggcag aaataagcat actcctgctt cccaaggcct gatgcttctc 19860
tccttccctc ctctcctcct ccttctcttc actctttctc tgcacacatg gaagaatggc 19920
tgccaggcat tgcccatttg gaaaagtaca gctcaatgga tatgaatcag cttgggcagg 19980
cgagaaatga ttcacgtctg accaaatcga tttagttcag gttgcccgtt ctgcatcttt 20040
tttcccttgc aattaaatga tgattgggtc tgatgggtgg aaggaagaga cagaatttaa 20100
tttgtttgccc tttgtagaaa gctgggggaca gcacagataa gggaagatgt ctcccatttg 20160

```


gcaaataact gatgcgagg tggagtggca gtggtgatgg ggatgctggt gccttcaggg 20220
cttctggggc gggcagtgca gctgggtggca gacggttcgg aactctacca tgttcccatc 20280
tgaaaactgt ggctgatcat gccactcct gaccttgctc cagggagtag acaaagacgt 20340
aagcttaatt aaccaccag acgtagctct tgaatccctg ggcatagtag ctgggtatag 20400
ttagagtttg ggagaggcat ggtcagcaaa acaacctccc tcatctctct gttgtcactc 20460
agagtcaagc tggctgctgc tgggtggtgct gacttctctt gctgcagatt tctccaatat 20520
gtttctgccc tgcacgcatt tgccaaatcc ctteggtttc ttgtgtctcg tggcagctta 20580
gctcctccag cccttggtatg aagaagcgtg ggaactcttt gcttcctttc cctcccgcag 20640
tgacatgcca tgccatgcca ctgcctcttc atctggctct atgacagtca ctcataagca 20700
cccgcatgta cccggccctg cactagctca tgacagctgc agtcaattgg gccaggtgct 20760
gtatctcatc cggcctctct agcaaccctc tgagatactg gtaatgtccc tgatgaagat 20820
atttactgag gcagaaatgg acgctcagtg aagcaagggt cctgatgtta tagcaatgag 20880
ctatgagtgg ccagagggag gagataagct caggcctgac accaaagccc atgctccttc 20940
tagtcaacca cagtgcctcc tatggtgaat gagttagtca gcaaccaaga cgcatgaggc 21000
cttctttttg gtgagccttg gctgggtgct gaggcttcag gtacaatcat gggttggaag 21060
agccctcctc tctctccaca gtctggcact atgaccctt ctggttatta acaaggcaaa 21120
gagagagagg gaagaaagca ggcaaataat gtgggttgct attcctagag attagaattt 21180
caggaaggat aaacacagcg ttctctccag aagtataaat aggaagactt cacacatgac 21240
tagaacgaga catgttttaa gtctgtcgag taaggcagtg atgaagtaga tttcccaga 21300
ttcactctcc ctctctggg tccccaggg ctttacttg tggcaacttt cagctcaggg 21360
agggaggaaa gcccctttca aagcttcaga tacttcctta aggtcagttt ctgcttaaag 21420
aaggccttta cattacttca tccctttgcc aaattaaact gaaaggaaac ctttcaagt 21480
tgattgctg gccctttcct gttcatttct cgtgggtacg ctttctaact ttctttcttt 21540
cttcccttct tcaggtgttg actttaagat gaagaccata gaggtagacg gcatcaaagt 21600
gcggatacag atctggtgag ctggggagga ggaggaggca gatgtaggag aagaggactt 21660
ctggctgctc cttagctgcc cctgccatgt gtaaaattcc taggcttcac ctgggataac 21720
tggccacctc tctgatggat ggaagcgaag tctcagaagc ccactctctc ctataagcct 21780
taatctccaa cctctaagaa actttagggg attgactaca agcaccaaag ggcaggaatt 21840
agaaggaaact ggcacactaa ccattgtgaa tttatctcag gattaggctt tgcccttggg 21900
ctgtgccaca ctatgttaag attggaagga aggaggctac acccccatc atttagggcg 21960
agaccctgag agagttcctc aggatagcat gatgaagttt ccacagtagc agaggggtgct 22020
gctgtggctc tctgcctgag gtcttggaag cactgccttt gccagggttt agagctccct 22080
ctcaattcca cagcagtagt ggcactgcct tcagagggtcc catagggact aggggtgtag 22140
cagcatcccc tgccaactcc catccaacca aatctggcca cagtggccag attccagaga 22200
gctgtccaag gcctgttctg gctgtggctt ctggtttctg ccaggaggggc agttggcagg 22260
aggggccaaag gccctgcagg cctggctcagc accagcacag atgaccaggc ctctgactgc 22320
agatccctgt ggggatccaa gcatccctgg tttttcacc ctttagctccc cagtttttcc 22380
tacaagggga cagctctgct cttcccctcc ccgtctgttc ccattggtccc tgctcctctg 22440
agggactggc tttctcctgc agggacactg cagggcagga gagataccag accatcaca 22500
agcagtacta tcggcggggc caggtaaagc accacattgg gggtttcaa gtgggaagct 22560
gccaccaca ctcccagctc tgggtatttg agatgtctgt ggcacggatc ccctaaatac 22620
agttcgctcgt cttggaggag cgcaggcgct ctttcagctg ttactgatc atttgtccgt 22680
ccattgttca tggcccactc actgcaggca ggcccctgcc ctaccccctg acttccacc 22740
tccatcctgg gtcaaagatc caggtcaaag catgtggtgt cttcctgctg tagagagttc 22800
tgtgatgggc ctgggaggcg gcagtgggtg ggtctgagag aagagatatt tctggatgct 22860
gagcaggagg aatgggagag tgggacccaa cctttaagtt tccacggccc cttctggccc 22920
catgactgca ctctctctgt gcatatcaca tctctctatt tctctctctc tcaggggata 22980
tttttggctc atgacattag cagcgagcgc tcttaccagc acatcatgaa gtgggtcagt 23040
gacgtggatg aggtaggaga tgccacctca ctgcccgggt gtggagaggg tgcctcaccg 23100
gggaaggcaa ggcgaggggc agatgggaag gcaaagtgtt ccagggaagct ttgccttcca 23160
cagccctgga tgaagacctc tgggtgagta agacatgggg aagaaaccga agctgccatg 23220
ccctcactct ctataacctg ccaggcctcc acggctgtgt ctttcccgga aatgaattag 23280
ttccaagtct tccctgtgag cagcttcttt cctgaaatct tgggaccagg tggagttgca 23340
agattgggat ctagtccctg ctctgcacaa tagctgtgga gccttgggaa gccatttgaa 23400
tctctgggt cccagttcc tgtagaatga gggctggact tacatccaat gtcctttcca 23460
gctctgatac cagtgttcta acccaaggaa gcaccagtct tagccagagt gtcttctacc 23520
ctaagctctc cccgtgatac ccttgaggtc agccatggca cttgggggag cctggcacct 23580

gcatccagtc ggcccaccct gtccttaggg ctctggaatt ggtggtgggc tggaggcagt 23640
 gcagactctg tagggaaaat tgggggggca ggcagcactc actggctgtt ctgcccattc 23700
 tttgtcccta gtacgcacca gaaggcgtcc agaagatcct tattgggaat aaggctgatg 23760
 aggagcagaa acggcagggt ggaagagagc aagggcagca ggtaagtggg gggaaaaggc 23820
 aagtccaccc caggctcctc gctgggcctc cagggccagt cctgagcgtg gggacctagg 23880
 ggtgtgttcc ccagtggcag gtcctccac acgtcccag caccccaagg ccctggggga 23940
 gtggccatcc tcggaaggct tgttgtctgg gtttcaggac agaagcccag agattcgggg 24000
 tccatccaga aacaaagacg tcataggcag caactctccc aagtccaggt ccccaaatgc 24060
 aggattgccc tctgtttaag agatcatccc cgtgttagta atgaaggact tcaagttgtc 24120
 aacctcttct ctgacagcat ccaggcctag ctgccatgtt acggtcgcga aatgatctcc 24180
 catcccaccc aacactcccc cactcctgtc cttcttacc aggaaagagc caggagggca 24240
 aatgaggaga caaagagcca cagctggaga agccatgggg gcagaaaagg taggaggatg 24300
 acgtgagggt aatgtccaag catgcaggga gaccatcctc ccagagagca gaaagaaata 24360
 ttggttattt ttttttctt tctttctttt tttttttttt tttgagatgg agtctcgctc 24420
 tgtcaccag gctagagtgc agtggcgcca tctcggtca ctgcaacctc tgctcctga 24480
 gttcaagcaa ttcttctgcc tcagcctccc aagtagctga gattacaggt gcatgccacc 24540
 acgcttggtt aatttttttg tatttttagt agagatgggg ttttgccata ttggccaggc 24600
 cggctctgaa ctccctaacct caggatgatc acctgcctca gtctcccaa gtgctgggat 24660
 tacaggcgtg agccactgtg cccagccaag attggtattt ctgagataag ttatccactc 24720
 agtccgtgga cctcaagagt tttcctctcc cttttcagtc aatagcgttc cattagtact 24780
 taaaatgaaa ttgattgttt ggtataaaat ataagacatg gtcattgacc aatttgaaag 24840
 tagaggcaaa gcctactagg atagtattta ttgagcactc tatgtgtggc actgtgctaa 24900
 ggcaagcgtt tttaagtga cgacccact gaatcatccc acaaccatgg atgggagaca 24960
 cactcagtct cctttaacag aagataaagc tggggcctac agagaatgta caacttgtcc 25020
 aaggtcacac agctagccat cagtggcagt gctgctattc aggtctggga ctgtgggact 25080
 ccagagccca tgttttttac gaggatgcca tactgccaca atggatgggt tctttatctc 25140
 ctgatatatg attgtgtgtt gggaggcgtg ggggtggcagc tggagaatg gagaggcata 25200
 tttgtggagg atcttcccc attctctgct acctctctt ggagctccca gtcccatctg 25260
 agaaattatc tactctgaga aatcgtcaca acacagcatg gttgtgagtg cagtggcaga 25320
 agcctgtgcc tggttgtatg ggccctctcc ctgccttact gactctcttt cagaaatgtc 25380
 cttctcttgc agctggcgaa ggagtatggc atggacttct atgaaacaag tgcctgcacc 25440
 aacctcaaca ttaaagaggt gagagccctg gtgaccaggc gcccgcctc tcgggctgag 25500
 tccagcagag gtgggaggag gagccataag atggacctta tccctcaggc cgctgcaggg 25560
 ttgccagggt agaggaggag acaactggact aacctgtgcc ctttggtttc cagtattca 25620
 cgcgtctgac agagctggtg ctgcaggccc ataggaagga gctggaaggc ctccggatgc 25680
 gtgccagcaa tgagtgggca ctggcagagc tggaggagga ggagggcaaa cccgagggac 25740
 cagcgaactc ttcgaaaacc tgctggtgct gactcctgtg tggggcacc caccagacac 25800
 ccctcttccc tcaggaggcc cgtgggcaga caggggagcc ggggctttgc cctgctgctg 25860
 tctctctgtg tgatgacct attgagtatc agtagccact actccccctg cctggcctg 25920
 agagcggctc tgetgtcatc tcaagcagcc cctgtcccca gcccgtccac cctggagtgg 25980
 tcttcttcag cctgtttccc cagccacagg cctgctacga ccccccacgat gtgccgcaag 26040
 cactgtctca ccatcccga cccaccagac aacagccagg gctggagtcc aggccacttt 26100
 cagctgctcc tttctccgtg categtgtct cttctctgct ttttctctct tccccactt 26160
 ctctttctct gacctctcc ctccggtgct tttcgtatca aagctcctca aacctcgctc 26220
 cccgtgtgtc ctgctgtgtg cagctcgctc tttccttct tcctaagcta tccaagggga 26280
 tggaccaggt ctctgtggga ggttccacc ttggatccag gaagaaccct ccacctgcc 26340
 tcgtgggtgg gccaaaggct acagggtgct tcttctctt ccccccacc cactgtccct 26400
 catgtgccat gggcctgct cccagtgac ctgcgaaagt ggagcatcga ggtaggaggg 26460
 aaacggcaac caggaggtcc tcgagcctgg ggctgcccta cctctaccca tccccgacc 26520
 agagctttgc ccttgcctgg ctgcccgcct gcctctttgg ggaactgagc tcagaggcag 26580
 gtgcttcaga gaaggaaaca aaatgagggg tggcagggat aaaaagtcac ctccattctc 26640
 tacctccat gcagcatgaa cacaatttct ctccacctgg ctcccaaatt taaagatgtg 26700
 gaccaaggcc tgtgggtact ccaggggcaa ggagagccct ggggtcagtg aactgtcag 26760
 gccaacatg cactccaca aggggagcat ttggaaatga aggactagct cctatgtatc 26820
 aggttaagag caaggagag ctggccaggg acagcagttt gcacagcaga ggggaatgta 26880
 gcaacagcag ggctcctag gcccactctt ccatttctta ggtaagaaga gcatttctc 26940
 agactccag gggaggact gagcctagcc ttcagcaacc aaggttctcc tgggacccaa 27000

```

agtttatggg agaagggcaa agacttcatg ggaagagaga aggaaggccc tgggtagaaa 27060
cgcttggtgc tgttctcttt ggcctttaag acaaagcgct catcttgccc tctacctcct 27120
gataggcttg aggggttgcc aaccacactg tggctacagg tggagggaaag aggactcctt 27180
cctccagagt gctatgttca ggaagtttct ttaaccccat atggcccaag agtagctcgt 27240
aggaggccct ttaaagacgg aacaagtaat ttaccagttc tactgggggtt cctgcccacc 27300
gtcccaaggt gggcgaggcc taggaagagg gtcattctta agccacacat tagctgcact 27360
gcgtggctgc agccaaaaca aagaactggg tgttgagtat tcatcaacta agaaccaaaa 27420
tccagggcac tcatatgtga aggataagaa cctcacttcc ttactcctcc aaaaagaagt 27480
ggggaaagaa ccatcaaacc tttctctctg acttaccaa ccaggaaaac agcaggagag 27540
gggtggctcag gacttaggga cagggtatag cttagatggt ggaaagcaaa ggagagcagg 27600
aagttgtaaa tctactggcta atgagaaaag gagacagcta actctaggat gaagctgtga 27660
ctaggctgga gttgcttctt tgaagatggg actccttggg tatcaagacc tatgccacat 27720
cacactgggg ctaggggaagt aggtgatgcc agccctcaag tctgtcttca gccagggact 27780
tgagaagtta tattgggcag tggctccaat ctgtggacca gtatttcagc tttccctgaa 27840
gatcaggcag ggtgccattc attgtctttc tctcctagcc ccctcaggaa agaaggacta 27900
tatttgtagt gtaccctagg ggttctggaa gggaaaacat ggaatcagga ttctatagac 27960
tgataggccc tatccacaag ggccatgact gggaaaaggt atgggagcag aaggagaatt 28020
gggatttttag ggtgcagcta cgctcaccct aaacttttgg tggcctgggg catgtcttga 28080
ggcccagact gttaaccagg ctctgctggc ctgtttactc gtcaccacct ctgcacctgc 28140
tgtcttgaga ctccatccag ccccaggcac gccacctgct cctgagcctc cactatctcc 28200
ctgtgacggg tgaacttcgt gtactgtgtc tcgggtccat atatgaattg tgagcagggt 28260
tcatctattt taaacacaga tgtttacaaa ataaagatta tttcaaacca cgggtgtggc 28320
tgcttgatg agtcttggg ggtaggtctc actcagacc tggcagtgat gtgggagggg 28380
gagaggcagt gctggtagaa gcagctccag aagcaaaggc aacagcagta gagtgaccac 28440
ggaagcggga aacattgtct tcccttctct accttcccta gtgccacctg cagggaggcc 28500
caaagcaag ccccgttgcc ctgcattggg ctggcactgc agaaataaga tgaacacag 28560
ttatcgagag gatgctgaac atctatgagc aggttttaaa gccaagatga gtctcatctg 28620
tttgtgtggg tcaggaacgg gtcttcttga aggcattgagg tgggactgga taatctttca 28680
gatttgtagt tggatacctc gggggagcag aggcagactg ggatctcagg actgcaggta 28740
tttcatactt tgggatatgg aattgatgga 28770

```

<210> 4

<211> 212

<212> PRT

<213> Rattus norvegicus

<400> 4

```

Met Ala Lys Gln Tyr Asp Val Leu Phe Arg Leu Leu Leu Ile Gly Asp
 1          5          10          15
Ser Gly Val Gly Lys Thr Cys Leu Leu Cys Arg Phe Thr Asp Asn Glu
 20          25          30
Phe His Ser Ser His Ile Ser Thr Ile Gly Val Asp Phe Lys Met Lys
 35          40          45
Thr Ile Glu Val Asp Gly Ile Lys Val Arg Ile Gln Ile Trp Asp Thr
 50          55          60
Ala Gly Gln Glu Arg Tyr Gln Thr Ile Thr Lys Gln Tyr Tyr Arg Arg
 65          70          75          80
Ala Gln Gly Ile Phe Leu Val Tyr Asp Ile Ser Ser Glu Arg Ser Tyr
 85          90          95
Gln His Ile Met Lys Trp Val Ser Asp Val Asp Glu Tyr Ala Pro Glu
100          105          110
Gly Val Gln Lys Ile Leu Ile Gly Asn Lys Ala Asp Glu Glu Gln Lys
115          120          125
Arg Gln Val Gly Arg Glu Gln Gly Gln Gln Leu Ala Lys Glu Tyr Gly
130          135          140
Met Asp Phe Tyr Glu Thr Ser Ala Cys Thr Asn Leu Asn Ile Lys Glu
145          150          155          160

```

Ser Phe Thr Arg Leu Thr Glu Leu Val Leu Gln Ala His Arg Lys Glu
 165 170 175
 Leu Asp Gly Leu Arg Thr Cys Ala Ser Asn Glu Leu Ala Leu Ala Glu
 180 185 190
 Leu Glu Glu Asp Glu Gly Lys Thr Glu Gly Pro Ala Asn Ser Ser Lys
 195 200 205
 Thr Cys Trp Cys
 210

<210> 5
 <211> 218
 <212> PRT
 <213> Human

<400> 5
 Met Ala Lys Gln Tyr Asp Val Leu Phe Arg Leu Leu Leu Ile Gly Asp
 1 5 10 15
 Ser Gly Val Gly Lys Thr Cys Leu Leu Cys Arg Phe Thr Asp Asn Glu
 20 25 30
 Phe His Ser Ser His Ile Ser Thr Ile Gly Val Asp Phe Lys Met Lys
 35 40 45
 Thr Ile Glu Val Asp Gly Ile Lys Val Arg Ile Gln Ile Trp Asp Thr
 50 55 60
 Ala Gly Gln Glu Arg Tyr Gln Thr Ile Thr Lys Gln Tyr Tyr Arg Arg
 65 70 75 80
 Ala Gln Gly Ile Phe Leu Val Tyr Asp Ile Ser Ser Glu Arg Ser Tyr
 85 90 95
 Gln His Ile Met Lys Trp Val Ser Asp Val Asp Glu Tyr Ala Pro Glu
 100 105 110
 Gly Val Gln Lys Ile Leu Ile Gly Asn Lys Ala Asp Glu Glu Gln Lys
 115 120 125
 Arg Gln Val Gly Arg Glu Gln Gly Gln Gln Lys Cys Pro Ser Leu Gln
 130 135 140
 Leu Ala Lys Glu Tyr Gly Met Asp Phe Tyr Glu Thr Ser Ala Cys Thr
 145 150 155 160
 Asn Leu Asn Ile Lys Glu Ser Phe Thr Arg Leu Thr Glu Leu Val Leu
 165 170 175
 Gln Ala His Arg Lys Glu Leu Glu Gly Leu Arg Met Arg Ala Ser Asn
 180 185 190
 Glu Leu Ala Leu Ala Glu Leu Glu Glu Glu Gly Lys Pro Glu Gly
 195 200 205
 Pro Ala Asn Ser Ser Lys Thr Cys Trp Cys
 210 215